

Emiliya V. Nosova,^{ab*} Olga A. Batanova,^a
 Nataliya N. Mochulskaya,^a Galina N. Lipunova^b

^a Ural Federal University

620002, 19 Mira St., Ekaterinburg, Russia

^b I. Ya. Postovskii Institute of Organic Synthesis,

Ural Branch of Russian Academy of Sciences,

620990, 22 Kovalevskaya St. / 20 Akademicheskaya St., Ekaterinburg, Russia

*email: emilia.nosova@yandex.ru

Synthetic approaches to 2-aryl/hetaryl- and 2-(hetaryl)ylidene derivatives of fluorinated 1,3-benzothiazin-4-ones

A series of 2-hetaryl- and 2-(hetaryl)ylidene substituted 5-fluoro-8-nitro-1,3-benzothiazin-4-ones was synthesized by interaction of 2,6-difluoro-3-nitrobenzoylisothiocyanate with C-nucleophiles. Cyclocondensation of poly-fluorobenzoylchlorides with aryl and hetaryl thioamides represents new approach to 1,3-benzothiazin-4-ones. Some compounds proved to be promising for further development of tuberculostatic agents.

Keywords: 1,3-benzothiazin-4-ones; 2-fluorobenzoylchloride; 2-fluorobenzoyl-isothiocyanate; indole; pyrrole; cyanomethylbenzimidazole; benzoylmethylbenzimidazole; thioamide; cyclocondensation; tuberculostatics.

Received: 28.06.2020. Accepted: 25.08.2020. Published: 07.10.2020.

© Emiliya V. Nosova, Olga A. Batanova, Nataliya N. Mochulskaya, Galina N. Lipunova, 2020

Introduction

Many synthetic benzothiazines are biologically actives, which play an important role in treatment of various diseases. Some 2-amino substituted 1,3-benzothiazin-4-ones (2-amino-1,3-benzothiazinones) represent a promising new class of antitubercular agents [1]. Other 1,3-benzothiazin-4-one derivatives, mostly 2-aryl and 2-(pyridin-2-yl) ones, are attractive due to their ability to suppress an oxidative stress-induced cardiomyocyte apoptosis [2]. Synthetic approaches to 2-amino-1,3-benzothiazin-4-ones are sufficiently developed, whereas not many ones are available for incorporation of C–C bond into position 2

[3]. The synthetic methods are limited to the following:

- interaction of 2-mercaptobenzoic acids with aryl/hetaryl nitriles [2];
- rearrangement of N-arylthiomethylaroylamides catalyzed by phosphorus oxychloride, followed by oxidation of 4*H*-1,3-benzothiazines with potassium permanganate [4];
- addition of C-nucleophiles to 2-fluorobenzoylisothiocyanates and subsequent intramolecular condensation [5].

The last approach opens wide opportunities for modification of position 2 of 1,3-benzothiazin-4-ones. Previously we

studied the interaction of polyfluorobenzo-ylisothiocyanates with CH-reactive benzimidazoles [5], 2,6-difluorobenzoylisothiocyanate with the same benzimidazoles and 2-cyanomethylpyridine [6], *o*-fluorobenzoylisothiocyanates with *N*-methylpyrrole and *N*-methylindole [7]. We presented only one example of 2-indolyl-5-fluoro-

8-nitro-1,3-benzothiazin-4-one in recent paper [8].

In this article, we wish to report new data on 2-substituted 5-fluoro-8-nitro-1,3-benzothiazinones and introduce efficient synthetic approach to 2-aryl/hetaryl derivatives of 1,3-benzothiazin-4-ones based on cyclocondensation of polyfluorobenzoyl chlorides with thioamides.

Experimental

^1H and ^{19}F NMR (nuclear magnetic resonance) spectra were recorded in dimethylsulfoxide- d_6 (DMSO- d_6) on the spectrometer "Bruker-Avance-400" (400 MHz), using tetramethylsilane as internal reference for ^1H NMR and CFCl_3 for ^{19}F NMR. Mass spectra were recorded on a SHIMADZU GCMS-QP2010 Ultra instrument with electron impact ionization (EI) of the sample. Microanalyses (C, H, N) were performed using the Perkin — Elmer 2400 elemental analyzer. Melting points were measured on the Stuart melting point apparatus SMP10 (AC/DC input 230 V AC, Merck supplier).

2,6-Difluorobenzoic acid **1**, 2,3,4,5-tetrafluorobenzoyl chloride **10a** and pentafluorobenzoyl chloride **10b** were purchased from Merck (CAS numbers 385-00-2, 94695-48-4, 2251-50-5). 3-Nitro-2,6-difluorobenzoic acid **2** was synthesized according to the literature [9]. Procedure for toluene solution of 2,6-difluoro-3-nitrobenzoylchloride **3** was reported [8]. 2-Benzoylmethylbenzimidazole was prepared from 2-methylbenzimidazole [5], 2-cyanomethyl-benzimidazole was synthesized by condensation of ethyl cyanoacetate with *o*-phenylenediamine [10]. Thioamides **11** were prepared by the addition of hydrogen sulfide to the corresponding nitriles [11].

5-Fluoro-8-nitro-2-(1-methylpyrrol-2-yl) — 1,3-benzothiazin-4-one (5). The solution of ammonium isothiocyanate (0.4758 g, 6.26 mmol) in acetonitrile (10 mL) was added to toluene solution of 2,6-difluoro-3-nitrobenzoylchloride **3** (0.88 mL, 6.26 mmol). Reaction mixture was stirred at 40 °C for 5 min, the precipitate of NH_4Cl was filtered off and 1-methylpyrrole (0.761 g, 9.39 mmol) was added to a solution of 2,6-difluoro-3-nitrobenzoylisothiocyanate **3**. The mixture was stirred at room temperature for 3 h, the precipitate of benzothiazinone **5** was filtered off and washed with hot ethanol (10 mL). Yield 1.72 g (90%), mp 194–196 °C. ^1H NMR, δ (ppm), J (Hz): 4.09 s (3H, CH_3), 6.30 dd (1H, H^4 , $^3J_{\text{HH}}$ 4.1, $^4J_{\text{HH}}$ 2.3), 7.29 dd (1H, H^3 , $^3J_{\text{HH}}$ 4.2, $^4J_{\text{HH}}$ 1.3), 7.37 m (1H, H^5), 7.62 dd (1H, H^6 , $^3J_{\text{HH}}$ 9.2, $^3J_{\text{FH}}$ 9.4), 8.70 dd (1H, H^7 , $^3J_{\text{HH}}$ 9.2, $^4J_{\text{FH}}$ 4.5). ^{19}F NMR, δ (ppm), J (Hz): — 99.05 dd (1F, $^3J_{\text{FH}}$ 9.5, $^4J_{\text{FH}}$ 4.0). MS (EI), m/z (I_{rel} (%)): 305 $[\text{M}]^+$ (14), 106 (100), 105 (15). Found, %: C 51.17; H 2.60; N 13.74. $\text{C}_{13}\text{H}_8\text{FN}_3\text{O}_3\text{S}$. Calculated, %: C 51.15; H 2.64; N 13.76.

Compounds **6–9** were synthesized by the same method.

5-Fluoro-8-nitro-2-(5-methoxy-1-methylindol-3-yl) — 1,3-benzothiazin-4-one (6). Yield 80%, mp 274–276 °C. ^1H NMR, δ (ppm), J (Hz): 3.83 s (3H, NCH_3), 3.93 s (3H, OCH_3), 7.01 d (1H, H^6 , $^3J_{\text{HH}}$

8.8), 7.54 d (1H, H⁷, ³J_{HH} 8.8), 7.65 dd (1H, H⁶, ³J_{HH} 9.3, ³J_{FH} 9.5), 7.97 s (1H, H⁴), 8.66 s (1H, H²), 8.70 dd (1H, H⁷, ³J_{HH} 9.3, ⁴J_{FH} 4.3). ¹⁹F{¹H} NMR, δ (ppm): — 99.27 s. MS (EI), *m/z* (I_{rel} (%)): 385 [M]⁺ (31), 187 (12), 186 (100), 171 (41), 143 (28). Found, %: C 56.13; H 3.15; N 10.87. C₁₈H₁₂FN₃O₄S. Calculated, %: C 56.10; H 3.14; N 10.90.

5-Fluoro-8-nitro-2-(2-methylindol-3-yl) — 1,3-benzothiazin-4-one (7). Yield 91%, mp 207–209 °C. ¹H NMR, δ (ppm), *J* (Hz): 2.87 s (3H, CH₃), 7.25 m (2H, C₆H₄), 7.45 m (1H, C₆H₄), 7.66 dd (1H, H⁶, ³J_{HH} 9.3, ³J_{FH} 9.6), 8.34 m (1H, C₆H₄), 8.71 dd (1H, H⁷, ³J_{HH} 9.3, ⁴J_{FH} 4.6), 12.5 br. s (1H, NH). ¹⁹F{¹H} NMR, δ (ppm): — 98.42 s. MS (EI), *m/z* (I_{rel} (%)): 355 [M]⁺ (21), 157 (12), 156 (100), 155 (50), 81 (10). Found, %: C 57.43; H 2.82; N 11.85. C₁₇H₁₀FN₃O₃S. Calculated, %: C 57.46; H 2.84; N 11.83.

1-(1,3-Dihydrobenzimidazol-2-yliden) — 1-(5-fluoro-8-nitro-4-oxo-4H-1,3-benzothiazin-2-yl) — acetonitrile (8). Yield 75%, mp 319–321 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.30 m (2H, C₆H₄), 7.50 dd (1H, H⁶, ³J_{HH} 9.2, ³J_{FH} 9.8), 7.64 m (2H, C₆H₄), 8.62 dd (1H, H⁷, ³J_{HH} 9.2, ⁴J_{FH} 4.4), 13.31 br. s (2H, NH). ¹⁹F{¹H} NMR, δ (ppm): — 97.49 s. MS (EI), *m/z* (I_{rel} (%)): 381 [M]⁺ (39), 183 (13), 182 (100), 155 (12), 103 (23), 81 (12). Found, %: C 53.50; H 2.15; N 18.36. C₁₇H₈FN₃O₃S. Calculated, %: C 53.54; H 2.11; N 18.37.

2-(1,3-Dihydrobenzimidazol-2-yliden) — 2-(5-fluoro-8-nitro-4-oxo-4H-1,3-benzothiazin-2-yl) — acetophenone (9). Yield 89%, mp 265–267 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.36 m (5H, C₆H₅), 7.53 m (4H, C₆H₄), 7.60 dd (1H, H⁶, ³J_{HH} 9.2, ³J_{FH} 9.5), 8.56 dd (1H, H⁷, ³J_{HH} 9.2, ⁴J_{FH} 4.5), 13.36 br. s (2H, NH). ¹⁹F{¹H} NMR, δ (ppm): — 98.39 s. MS (EI), *m/z* (I_{rel} (%)): 460 [M]⁺ (24), 432 (13), 431 (41), 355 (25), 261 (46), 260 (100), 206 (16), 156 (16), 105

(45), 77 (71), 51 (10). Found, %: C 60.03; H 2.83; N 12.20. C₂₃H₁₃FN₄O₄S. Calculated, %: C 60.00; H 2.85; N 12.17.

6,7,8-Trifluoro-2-phenyl-1,3-benzothiazin-4-one (12a). Tetrafluorobenzoylchloride **10a** (0.85 g, 4 mmol) was added to thiobenzamide **11a** (0.397 g, 2.9 mmol) in dry toluene (8 mL), reaction mixture was refluxed for 3 h and then cooled. The precipitate of benzothiazinone **12a** was filtered off and recrystallized from DMSO. Yield 0.646 g (76%), mp 160–162 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.62 m (2H, Ph), 7.73 m (1H, Ph), 8.15 ddd (1H, H⁵, ³J 10.3, ⁴J 7.4, ⁵J 2.2), 8.19 m (2H, Ph). ¹⁹F NMR, δ (ppm), *J* (Hz): 151.84 ddd (1F, F⁷, ³J 22.5, ³J 21.5, ⁴J 7.4), 135.10 ddd (1F, F⁸, ³J 21.5, ⁴J 6.2, ⁵J 2.2), 132.50 ddd (1F, F⁶, ³J 22.5, ³J 10.3, ⁴J 6.2). MS (EI), *m/z* (I_{rel} (%)): 293 [M]⁺ (11), 190 (100), 162 (30). Found, %: C 57.51, H 1.88, N 4.62. C₁₄H₆F₃NOS. Calculated, %: C 57.34, H 2.06, N 4.78.

Compounds **12b-h** were synthesized by the same method.

6,7,8-Trifluoro-2-(*p*-chlorophenyl) — 1,3-benzothiazin-4-one (12b). Yield 59%, mp 204–206 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.73 d (2H, H^{3,5}, ³J 8.8), 8.23 d (2H, H^{2,6}, ³J 8.8), 8.24 ddd (1H, H⁵, ³J 10.6, ⁴J 7.5, ⁵J 2.1). ¹⁹F NMR, δ (ppm), *J* (Hz): 151.68 ddd (1F, F⁷, ³J 22.5, ³J 21.2, ⁴J 7.5), 135.02 ddd (1F, F⁸, ³J 21.5, ⁴J 6.3, ⁵J 2.1), 132.29 ddd (1F, F⁶, ³J 22.5, ³J 10.6, ⁴J 6.3). MS (EI), *m/z* (I_{rel} (%)): 327 [M]⁺ (4), 190 (100), 162 (27). Found, %: C 51.42, H 1.66, N 4.13. C₁₄H₅F₃NOSCl. Calculated, %: C 51.31, H 1.54, N 4.27.

6,7,8-Trifluoro-2-(*p*-tolyl) — 1,3-benzothiazin-4-one (12c). Yield 71%, mp 184–186 °C. ¹H NMR, δ (ppm), *J* (Hz): 2.46 s (3H, CH₃), 7.43 d (2H, H^{3,5}, ³J 8.0), 8.09 d (2H, H^{2,6}, ³J 8.0), 8.14 ddd (1H, H⁵, ³J 10.0, ⁴J 7.5, ⁵J 2.3). Found, %: C 58.75, H 2.70, N 4.41. C₁₅H₈F₃NOS. Calculated, %: C 58.63, H 2.62, N 4.56.

6,7,8-Trifluoro-2-(pyridyl-2) — 1,3-benzothiazin-4-one (12d). Yield 76%, mp 166–168 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.75 dd (1H, H⁵, ³*J* 8.0, ³*J* 4.0), 8.10 td (1H, H⁴, ³*J* 8.0, ⁴*J* 1.8), 8.13 ddd (1H, H⁵, ³*J* 10.4, ⁴*J* 7.4, ⁵*J* 2.2), 8.38 d (1H, H³, ³*J* 8.0), 8.79 dd (1H, H⁶, ³*J* 4.0, ⁴*J* 1.8). ¹⁹F NMR, δ (ppm), *J* (Hz): 151.95 ddd (1F, F⁷, ³*J* 22.5, ³*J* 21.1, ⁴*J* 7.4), 135.64 ddd (1F, F⁸, ³*J* 21.1, ⁴*J* 6.2, ⁵*J* 2.2), 132.70 ddd (1F, F⁶, ³*J* 22.5, ³*J* 10.4, ⁴*J* 6.2). Found, %: C 52.95, H 1.63, N 9.67. C₁₃H₅F₃N₂OS. Calculated, %: C 53.06, H 1.71, N 9.52.

5,6,7,8-Tetrafluoro-2-phenyl-1,3-benzothiazin-4-one (12e). Yield 80%, mp 165–167 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.63 m (2H, Ph), 7.76 m (1H, Ph), 8.17 m (2H, Ph). MS (EI), *m/z* (*I*_{rel} (%)): 311 [M]⁺ (7), 208 (100), 180 (24), 111 (5). Found, %: C 53.83, H 1.81, N 4.67. C₁₄H₅F₄NOS. Calculated, %: C 54.02, H 1.62, N 4.50.

Results and discussion

We developed an efficient synthetic approach to 2-hetaryl/(hetaryl)ylidene-substituted fluorinated 1,3-benzothiazinones, for this purpose we studied the interaction of the range of C-nucleophiles (indoles, N-methylpyrrole, 2-cyanomethyl- and 2-benzoylmethyl- benzimidazoles) with 2,6-difluoro-3-nitrobenzoylisothiocyanate **4** in acetonitrile at room temperature (Figure 1). According to ¹H and ¹⁹F NMR spectra, the reaction leads to the formation of 1,3-benzothiazin-4-ones **5–9**, the intermediate addition products were not isolated, and fluorine atom at C⁵ was not subjected to substitution with nucleophilic reagent. It is worth noting that the intramolecular cyclization proceeded at milder reaction conditions than in the case of 2,6-difluoro- and 2,3,4,5-tetrafluorobenzoyl derivatives (refluxing in acetonitrile or

5,6,7,8-Tetrafluoro-2-(*p*-chlorophenyl) — 1,3-benzothiazin-4-one (12f). Yield 62%, mp 186–188 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.66 d (2H, H^{3:5}, ³*J* 8.8), 8.20 d (2H, H^{2:6}, ³*J* 8.8). Found, %: C 48.83, N 3.97. C₁₄H₄F₄NOSCl. Calculated, %: C 48.64, H 1.17, N 4.05.

5,6,7,8-Tetrafluoro-2-(*p*-tolyl) — 1,3-benzothiazin-4-one (12g). Yield 74%, mp 191–193 °C. ¹H NMR, δ (ppm), *J* (Hz): 2.46 s (3H, CH₃), 7.43 d (2H, H^{3:5}, ³*J* 8.4), 8.08 d (2H, H^{2:6}, ³*J* 8.4). Found, %: C 55.46, H 2.24, N 4.19. C₁₅H₇F₄NOS. Calculated, %: C 55.39, H 2.17, N 4.31.

5,6,7,8-Tetrafluoro-2-(pyridyl-2) — 1,3-benzothiazin-4-one (12h). Yield 77%, mp 182–184 °C. ¹H NMR, δ (ppm), *J* (Hz): 7.76 ddd (1H, H⁵, ³*J* 8.0, ³*J* 4.8, ⁴*J* 0.8), 8.10 td (1H, H⁴, ³*J* 8.0, ⁴*J* 1.8), 8.36 dd (1H, H³, ³*J* 8.0, ⁴*J* 0.8), 8.79 dd (1H, H⁶, ³*J* 4.8, ⁴*J* 1.8). Found, %: C 49.92, N 9.09. C₁₃H₄F₄N₂OS. Calculated, %: C 50.01, H 1.29, N 8.97.

dimethylformamide in the presence of trimethylamine [7]).

The signals of H⁶ and H⁷ in ¹H NMR spectra of benzothiazinones **5–9** exhibit more complicated multiplicity than in the case of 2,5-diaminobenzothiazinones [8], which indicates that the fluorine atom remains in position 5. To prove that the alternative product of cyclization, 5-fluoro-6-nitro isomer, was not formed ¹⁹F NMR spectra without suppression of F-H spin-spin interaction were registered. In such spectra of compounds **5–9** double doublet signals with ³*J*_{FH} = 9.5–10.1 Hz and ⁴*J*_{FH} = 3.9–4.0 Hz are present, so the formation of 5-fluoro-8-nitroisomers was confirmed. The peaks of molecular ions in the mass spectra of benzothiazinones **5–9** have a relative intensity of 14–39%.

Thus, we demonstrated the difference in behavior of C-nucleophiles and N-nucleophiles under the reaction with 2,6-difluoro-3-nitrobenzoylisothiocyanate **4**: application of C-nucleophiles allows to obtain derivatives of 5-fluoro-8-nitrobenzothiazinone, whereas the reaction with cycloalkylimines fails to avoid the nucleophilic substitution of fluorine at position 5. The proposed strategy opens wide opportunities for varying the substituent at position 2 of 8-nitrobenzothiazin-4-ones.

We presented novel one-stage synthetic approach to 2-substituted fluorine-containing 1,3-benzothiazin-4-ones based on cyclocondensation of polyfluoroben-

zoyl chlorides with thioamides as S,N-dinucleophiles. New 6,7,8-trifluoro- and 5,6,7,8-tetrafluoro-derivatives of 1,3-benzothiazin-4-ones **12a-h** were obtained by the reaction of polyfluorobenzoyl chlorides **10a,b** and thioamides **11a-d** in boiling toluene for 3 h in 59–80% yields (Figure 2), notably that intermediate N-arylation products were not isolated. Signals of NH groups are absent in ^1H NMR spectra of compounds **12a-h**, spectra of 6,7,8-trifluorobenzothiazinones **12a-d** exhibited ddd signal of fluoroarene residue H^5 proton at 8.13–8.24 ppm. The number of signals in ^{19}F NMR spectra is in accordance with the structure of benzothiazinones **12**. The peaks of molecular ions in the mass




Fig. 1. Synthesis of 5-fluoro-8-nitro-1,3-benzothiazin-4-ones **5-9**

spectra of benzothiazinones **12** have low relative intensity of 4–11%. The ions m/z 190 or m/z 208 with 100% intensity were observed for benzothiazinones **12**, moreover, peaks m/z 162 or m/z 180 correspond to ions $[M-RCN-CO]^+$. The most abundant peak was reported to be characteristic for elimination of RCN fragment from molecular ions of 2-R-6,7,8-trifluoro-1,3-benzothiazin-4-ones [5–7].

The presented approach allows to obtain a variety of 2-aryl/hetaryl-substituted 1,3-benzothiazinones and successfully complements the previously described cy-

clocondensation of polyfluorobenzoylchlorides with benzimidazol-2-thiones as cyclic S,N-dinucleophiles leading to [b]-annelated fluorobenzothiazinones [12]. Unfortunately, we failed to obtain 5-fluoro- and 5-fluoro-8-nitro analogs using the method shown in Figure 2.


Tuberculostatic activity of polyfluorinated benzothiazinones **12** was studied at two laboratories, namely Ural Research Institute for Phthisiopulmonology (URIP) and University of Illinois, Chicago Institute for Tuberculosis Research (INR); the data are presented in Table 1.

Table 1

Data on tuberculostatic activity of fluorinated 2-aryl/pyridyl-1,3-benzothiazin-4-ones **12 against *Mycobacterium tuberculosis* $H_{37}R_v$ ***

Comp	MIC values (URIP data), $\mu\text{g/mL}$	% Inhibition at 128 $\mu\text{g/mL}$ (ITR data)		MIC values, $\mu\text{g/mL}$ (ITR data)		IC_{50} , $\mu\text{g/mL}$ (ITR data)
		MABA	LORA	MABA	LORA	
12b	12.5	28	66	-	-	—
12c	12.5	95	100	58.0	52.3	>128
12d	0.3	—	—	>128	>128	—
12f	3.12	0	100	—	3.7	53.5
12g	nd	74	99	—	26.3	—
12h	12.5	0	100	—	55.8	65.9

* MIC—Minimal inhibitory concentration; IC_{50} — the half maximal inhibitory concentration; URIP — Ural Research Institute for Phthisiopulmonology; INR — Chicago Institute for Tuberculosis Research; MABA — microplate Alamar Blue assay; LORA — low-oxygen recovery assay.



10: Y = H (**a**), F (**b**); **11**: R = Ph (**a**), 4-Cl-C₆H₄ (**b**), 4-Me-C₆H₄ (**c**), 2-Py (**d**); **12**: Y = H, R = Ph (**a**), 4-Cl-C₆H₄ (**b**), 4-Me-C₆H₄ (**c**), 2-Py (**d**); Y = F, R = Ph (**e**), 4-Cl-C₆H₄ (**f**), 4-Me-C₆H₄ (**g**), 2-Py (**h**).

Fig. 2. Synthesis of polyfluorinated 2-aryl/pyridyl-1,3-benzothiazin-4-ones **12a-h**

According to trials conducted in URIP, benzothiazinone **12d** exhibited the highest activity (MIC 0.3 µg/mL, isoniazide as reference compound with MIC 0.15 µg/mL). Data obtained in ITR revealed

12f as the most promising derivative towards the dormant multi-resistant strain of micobacteria H₃₇R_v-CA-luxAB (MIC 3.7 µg/mL, rifampicinum as reference compound with MIC 8.26 µg/mL).

Conclusions

To sum up, we developed efficient synthetic approaches to fluorine-containing 1,3-benzothiazin-4-ones bearing aryl, hetaryl and (hetaryl)ylidene residues at po-

sition 2 and demonstrate that some of them are promising for design of new antitubercular agents.

Acknowledgements

The work was carried out with financial support from the Ministry of Science and Higher Education of Russian Federation (project № FEUZ-2020-0058 (H687.42B.223/20)).

References

1. Chetty S, Ramesh M, Singh-Pillay A, Soliman M. Recent advancements in the development of anti-tuberculosis drugs. *Bioorg Med Chem Lett*. 2017;27(3):370–86. DOI: 10.1016/j.bmcl.2016.11.084
2. Kimura H, Sato Y, Tajima Y, Suzuki H, Yukitake H, Imaeda T, Kajino M, Oki H, Takizawa M, Tanida S. BTZO-1, a Cardioprotective Agent, Reveals that Macrophage Migration Inhibitory Factor Regulates ARE-Mediated Gene Expression. *Chem Biology*. 2010;17(12):1282–94. DOI: 10.1016/j.chembiol.2010.10.011
3. Nosova EV, Lipunova GN, Charushin VN, Chupakhin ON. Synthesis and biological activity of 2-amino- and 2-aryl(hetaryl) substituted 1,3-benzothiazin-4-ones. *Mini-Reviews in Medicinal Chemistry*. 2019;19(12):999–1014. DOI: 10.2174/1389557518666181015151801
4. Szabo J, Bani-Akoto E, Dombi G, Gunther G, Bernath G, Fodor L. Ring-closure reaction of N-arylthiomethylaroylamides to 1,3-benzothiazines. *J. Heterocyc. Chem*. 1992;29(5):1321–4. DOI:10.1002/jhet.5570290545
5. Nosova EV, Lipunova GN, Laeva AA, Charushin VN. Polyfluorobenzoyl chlorides and isothiocyanates in reactions with CH-reactive benzimidazoles. *Russ. Chem. Bull*. 2005;54(3):733–7. DOI: 10.1007/s11172-005-0312-6
6. Nosova EV, Laeva AA, Trashakhova TV, Golovchenko AV, Lipunova GN, Slepukhin PA, Charushin VN. Fluorine-containing heterocycles: XVIII. Monofluoro derivatives of quinazolines and 1,3-benzothiazin-4-ones. *Russ J Org Chem*. 2009;45(6):904–12. doi:10.1134/S1070428009060189

7. Nosova EV, Poteeva AD, Lipunova GN, Slepukhin PA, Charushin VN. Synthesis of Fluorine-Containing 2-Pyrrolyl- and 2-Indolyl-Substituted 1,3-Benzothiazin-4-ones. *Russ J Org Chem.* 2019;55(3):384–7.
DOI: 10.1134/S1070428019030205
8. Nosova EV, Batanova OA, Lipunova GN, Charushin VN. Synthesis of novel 8-nitro-substituted 1,3-benzothiazin-4-ones. *Mendeleev Communications.* 2020;30(4):427–9.
DOI: 10.1016/j.mencom.2020.07.007
9. Yoshida Y, Barrett D, Azami H, Morinaga C, Matsumoto S, Matsumoto Y, Takasugi H. Studies on Anti-Helicobacter pylori Agents. Part 1: Benzylxyisoquinoline Derivatives. *Bioorg Med Chem.* 1999;7(11):2647–66.
DOI: 10.1016/S0968-0896(99)00203–5
10. Mohareb RM, Abdallah AEM, Mohamed AA. Synthesis of Novel Thiophene, Thiazole and Coumarin Derivatives Based on Benzimidazole Nucleus and Their Cytotoxicity and Toxicity Evaluations. *Chem Pharm Bull.* 2018;66(3):309–18.
DOI: 10.1248/cpb.c17–00922
11. Mochul'skaya NN, Andreiko AA, Kodess MI, Vasileva EB, Filyakova V. I, Gubaidullin AT, Litvinov IA, Sinyashin OG, Aleksandrov GG, Charushin VN. Annulation of the thiazole ring to 1,2,4-triazines by tandem A_N-A_N or $S_N^H-S_N^H$ reactions. *Russ Chem Bull, Int Ed.* 2004;53(6):1279–89.
DOI: 10.1023/B:RUCB.0000042287.32992.54
12. Lipunova GN, Nosova EV, Mokrushina GA, Ogloblina EG, Aleksandrov GG, Charushin VN. Fluorocontaining Heterocycles: IX. Derivatives of Imidazo[2,1-*b*][1,3]benzothiazine. *Russ J Org Chem.* 2003;39(2):248–56.
DOI: 10.1023/A:1025548505109