






**2-(4-Oxo-1,3-thiazolidin-2-ylidene)acetamide as a promising
scaffold for designing new antifungal compounds**

Konstantin L. Obydenov* , Tatiana A. Kalinina , Daria V. Ryabova,
Maria F. Kosterina, Tatiana V. Glukhareva 

Institute of Chemical Engineering, Ural Federal University, Ekaterinburg
620009, Russia

* (E-mail: k.l.obydenov@urfu.ru)




Figure S1 The ^1H NMR spectrum (400 MHz, DMSO) of 3c.




Figure S2 The ^{13}C NMR spectrum (400 MHz, DMSO) of 3c.




Figure S3 The ^1H NMR spectrum (400 MHz, DMSO) of 4a.




Figure S4 The ^{13}C NMR spectrum (400 MHz, DMSO) of 4a.




Figure S5 The ^1H NMR spectrum (400 MHz, DMSO) of 4b.




Figure S6 The ^{13}C NMR spectrum (400 MHz, DMSO) of 4b.




Figure S7 The ^1H NMR spectrum (400 MHz, DMSO) of 4c.




Figure S8 The ^{13}C NMR spectrum (400 MHz, DMSO) of 4c.




Figure S9 The ¹H NMR spectrum (400 MHz, DMSO) of 4e.




Figure S10 The ¹³C NMR spectrum (400 MHz, DMSO) of 4e.